

CROSSCAP NUMBERS OF TORUS KNOTS

MASAKAZU TERAGAITO

ABSTRACT. The crosscap number of a knot in the 3-sphere is the minimal genus of non-orientable surface bounded by the knot. We determine the crosscap numbers of torus knots.

1. INTRODUCTION

For a knot K in the 3-sphere S^3 , the *crosscap number* of K , denoted by $c(K)$, is defined to be the minimal first betti number of compact, connected, non-orientable surfaces F with $\partial F = K$ [3]. (For the trivial knot, it is defined to be zero instead of one.) Since the crosscap number is an analogy of the genus of a knot, it is also called the non-orientable genus [12]. Clark [3] showed that a knot has crosscap number one if and only if it is a 2-cable of some knot. In [15], we showed that genus one, crosscap number two knots are doubled knots. Since a genus one knot has crosscap number at most three ([3, 11]), this determines the crosscap numbers of genus one knots. However it is hard to determine the crosscap number of a knot, in general. See also [1, 8, 11, 18].

Let F be a surface bounded by a knot K . Then it can be assumed that F meets the regular neighborhood $N(K)$ of K in a collar neighborhood of ∂F in F , which is an annulus. Then $F \cap \partial N(K)$ is an essential simple closed curve on $\partial N(K)$. Its (unoriented) isotopy class is referred to as the *boundary slope* of F , which is parameterized by integers in the usual way (see [13]). The boundary slope of F is uniquely determined by F . Clearly, if F is orientable, then the boundary slope is zero. But this is not the case when F is non-orientable. Then the boundary slope must be an even integer by homological reason (see [2]). If F is a spanning surface of K , that is $\partial F = K$, then a new non-orientable spanning surface F' of K is obtained by adding a small half-twisted band to F locally. The boundary slope of F' is (that of F) ± 2 . Thus any even integer can be the boundary slope of some non-orientable spanning surface of K . The first betti number of F is denoted by $\beta_1(F)$. In particular, $\beta_1(F)$ is equal to the genus of F when F is non-orientable.

In this paper, we determine the crosscap number of a torus knot K and the boundary slope of a non-orientable surface bounded by K , which realizes its crosscap number $c(K)$, simultaneously. In fact, we show that the boundary slope of a non-orientable spanning surface of K realizing $c(K)$ is uniquely determined by K . This does not hold in general. For example, the figure-eight knot has crosscap

number two, and it bounds two once-punctured Klein bottles with boundary slopes 4 and -4 , which are two checkerboard surfaces of a standard diagram. Another remarkable example is the $(-2, 3, 7)$ -pretzel knot, which bounds two once-punctured Klein bottles with boundary slopes 16 and 20. See [8]. For torus knots, the crosscap numbers are given by using a positive integer $N(x, y)$ introduced by Bredon and Wood [2], which is the minimal genus of closed, connected, non-orientable surfaces contained in the lens space $L(x, y)$. It is easy to calculate $N(x, y)$ by using continued fractions or a recursive formula (see Section 5).

Let T be a standard torus in S^3 . It decomposes S^3 into two solid tori V and W . Let $f : S^1 \times D^2 \rightarrow V$ be a homeomorphism. This determines the longitude-meridian system of V by setting $\lambda = f(S^1 \times *)$, $*$ $\in \partial D^2$, and $\mu = f(* \times \partial D^2)$, $*$ $\in S^1$, which gives a basis of $H_1(T)$. We assume that λ is a preferred longitude, that is, λ bounds a disk in W . Let $T(p, q)$ be the torus knot of type (p, q) lying on T , which represents $p[\lambda] + q[\mu]$ in $H_1(T)$. Note that p and q are coprime. Since all $T(p, q), T(-p, q), T(p, -q), T(-p, -q), T(q, p)$ are equivalent (there is a homeomorphism of S^3 sending one to the other), they have the same crosscap number. Therefore we always assume $p, q > 0$. The torus knot $T(p, q)$ is said to be *odd* (resp. *even*) if pq is odd (resp. even). Furthermore, if $T(p, q)$ is odd (resp. even), then we always assume that $p > q$ (resp. p is even).

Theorem 1.1. *Let K be the non-trivial torus knot of type (p, q) , where $p, q > 0$, and let F be a non-orientable spanning surface of K with $\beta_1(F) = c(K)$.*

- (1) *If K is even, then $c(K) = N(p, q)$ and the boundary slope of F is pq .*
- (2) *If K is odd, then $c(K) = N(pq - 1, p^2)$ (resp. $N(pq + 1, p^2)$) and the boundary slope of F is $pq - 1$ (resp. $pq + 1$) if $xq \equiv -1 \pmod{p}$ has an even (resp. odd) solution x satisfying $0 < x < p$.*

Remark that the equation $xq \equiv -1 \pmod{p}$ in (2) has the unique solution x satisfying $0 < x < p$. See Section 4.

In general, the crosscap number is not additive under the connected sum operation [11]. But we have:

Theorem 1.2. *If $K_1, K_2, \dots, K_n$ are torus knots, then*

$$c(K_1 \# K_2 \# \dots \# K_n) = c(K_1) + c(K_2) + \dots + c(K_n).$$

2. PRELIMINARIES

Let $K = T(p, q)$ be the non-trivial torus knot of type (p, q) , and let $E(K)$ be its exterior. Let F be a non-orientable surface bounded by K realizing its crosscap number. We may assume that $F \cap N(K)$ is an annulus, and $F \cap E(K)$ is also denoted by F . As noted in Section 1, ∂F ($\subset \partial E(K)$) determines the boundary slope r , which is an even integer.

Lemma 2.1. *F is incompressible in $E(K)$.*




FIGURE 1.

Proof. Assume that F is compressible in $E(K)$, and let D be a compressing disk for F . Note that ∂D is orientation-preserving in F .

If ∂D is separating in F , then compression along D gives two surfaces F_1 and F_2 , where $\partial F_1 = \partial F$ and F_2 is closed. Then F_2 is orientable, and hence F_1 is non-orientable. It is easy to see that $\beta_1(F_1) \leq \beta_1(F) - 2$. This contradicts the minimality of $\beta_1(F)$. Therefore ∂D is non-separating in F .

Let F' be the resulting surface obtained by compressing F along D . Since $\beta_1(F') = \beta_1(F) - 2$, F' must be orientable by the minimality of F . But if we add a small half-twisted band to F' (after extending F' to K radially in $N(K)$), then we obtain a non-orientable surface F'' bounded K with $\beta_1(F'') = \beta_1(F') + 1 < \beta_1(F)$. This contradicts the minimality of F . $\square$

Recall that K lies on the standard torus T . Let $A = T \cap E(K)$, and let a_1 and a_2 be the components of ∂A . Then it is well known that A is an essential (incompressible and boundary-incompressible) annulus in $E(K)$ and that each component a_i of ∂A has slope pq on $\partial E(K)$. By an isotopy of F , we may assume that F and A intersect transversely, and therefore $F \cap A$ consists of arcs and loops. Since F and A are incompressible, we can remove any loop component of $F \cap A$ bounding a disk on either F or A . Furthermore, we may assume that ∂F meets each a_i in the same direction (after giving them orientations). Then $F \cap A$ contains exactly $\Delta(r, pq) = |r - pq|$ arcs, since ∂F meets a_i in $\Delta(r, pq)$ points, where $\Delta(r, pq)$ denotes the minimal geometric intersection number between two slopes r and pq on $\partial N(K)$.

Lemma 2.2. *If $F \cap A$ contains an arc component α , then α connects distinct boundary components of A .*

Proof. Suppose that the ends of α lie in a_1 , say. Without loss of generality, we can assume that α is outermost on A . That is, α cuts off a disk D from A such that $\text{Int} D \cap F = \emptyset$. Let $\partial D = \alpha \cup \beta$, where β is a subarc of a_1 . See Figure 1.

By doing boundary-compression along D , we obtain a connected surface F' . Note that $\beta_1(F') = \beta_1(F) - 1$ by an Euler characteristic calculation. Since $\partial F'$ still meets a meridian of K in one point, F' can be extended to a spanning surface




FIGURE 2.

of K . Thus F' is orientable, because of the minimality of F . Then F is obtained from a Seifert surface F' of K by adding a half-twisted band locally. Since K has genus $(p-1)(q-1)/2$ ([13]), it implies that $c(K) = \beta_1(F) = (p-1)(q-1) + 1$. By [11, Proposition 1.3], $c(K) \leq \min\{(p-1)q/2, (q-1)p/2\}$. (A standard diagram of K has $(p-1)q$ or $(q-1)p$ crossings.) This gives a contradiction easily. $\square$

Thus, if $|pq - r| \neq 0$, then $F \cap A$ contains no loop component. If $|pq - r| = 0$, then $F \cap A$ may contain loop components which are essential in A . Since r is even, $|pq - r| = 0$ happens only when K is even. Also, if K is odd, then $F \cap A$ contains at least one arc.

We introduce an operation called a *disk splitting*. This operation will be used to determine the boundary slope of F in the following sections. If $F \cap A$ contains at least two arcs, then there are two arcs α and β of $F \cap A$, which cut a rectangle D from A such that $\text{Int}D \cap F = \emptyset$. See Figure 2. Let F' be the surface obtained by splitting F along D . That is, take a product neighborhood $D \times [0, 1]$ of D in $E(K)$, and let $F' = (F - D \times [0, 1]) \cup D \times \{0, 1\}$. Note that this disk splitting does not change Euler characteristic. (If F' is disconnected, then $\chi(F')$ is the sum of Euler characteristic of its components.)

3. THE CASE WHERE K IS EVEN

In this section, we prove Theorem 1.1(1).

Suppose that K is even. Recall that p is assumed to be even, and that r is the boundary slope of F .

Proposition 3.1. $r = pq$.

Proof. Suppose not. Let $\Delta = \Delta(r, pq) = |r - pq|$. Then Δ is even and $\Delta \geq 2$.

Claim 3.2. $p \geq 4$.

Proof of Claim 3.2. If $p = 2$, then $c(K) = 1$ and F is a Möbius band. Consider r -Dehn surgery $K(r)$. That is, $K(r)$ is the union of $E(K)$ and a solid torus J glued




FIGURE 3.

to $E(K)$ along their boundaries in such a way that r bounds a meridian disk in J . Thus $K(r)$ contains a projective plane, which is obtained by capping the Möbius band off by a meridian disk of J . Then $K(r)$ is P^3 or reducible. By [10], the former is impossible, and so $K(r)$ is reducible. Also pq is the only slope yielding a reducible manifold. Hence $p \geq 4$. $\square$

Now, $F \cap A$ consists of Δ arcs. There are mutually disjoint $\Delta/2$ rectangles on A cut by these arcs. We perform $\Delta/2$ disk splittings to obtain two surfaces F_1 and F_2 . Note that $\chi(F) = \chi(F_1) + \chi(F_2)$. One surface is connected and has a single boundary component, whose slope is pq , and the other has $\Delta/2$ boundary components, each of which is inessential on $\partial E(K)$. In particular, the latter does not contain a non-orientable component. See Figure 3 (where $\Delta = 6$). We can assume that F_1 has a single boundary component. Since the slope pq of ∂F_1 is not zero, F_1 must be non-orientable. If we show $\chi(F_2) < 0$, then $\chi(F) < \chi(F_1)$, which contradicts the minimality of F .

Number the points of $a_1 \cap \partial F$ $1, 2, \dots, \Delta$ along a_1 . This induces the labeling of the points of $a_2 \cap \partial F$ through the arcs of $F \cap A$. See Figure 3. The annulus A splits $E(K)$ into two solid tori U_1 and U_2 . We assume U_i contains F_i for $i = 1, 2$. Let $A_i = \partial E(K) \cap U_i$. Then for some permutation σ of $\{1, 2, \dots, \Delta\}$, the point with label i in a_1 is connected to one with label $\sigma(i)$ in a_2 by the arcs of $A_2 \cap \partial F$. (In Figure 3, $\sigma = (153)(264)$.) Note that σ has an even number of orbits, all of which have the same length $\ell(\sigma)$, say, by the definition. (Thus σ has $\Delta/\ell(\sigma)$ orbits.)

Since F_2 has $\Delta/2$ boundary components, and F_2 has no closed components, $|F_2| \leq \Delta/2$. ($|\dots|$ denotes the number of components.) In fact, the boundary components of F_2 belong to the same component of F_2 when they correspond to the same orbit of σ . Thus $|F_2| \leq \Delta/2\ell(\sigma)$.

Claim 3.3. F_2 contains no disk components.

Proof of Claim 3.3. Suppose not. We can regard F_2 as the union of $F \cap U_2$ and the rectangles on A used in the disk splittings. Let D be a disk component of F_2 . Then we have $\ell(\sigma) = 1$. We can take a loop f on ∂U_2 , which is the union of an arc of $F \cap A$ and an arc of $\partial F \cap A_2$ and which meets ∂D . Then this loop f bounds a disk in D . Since f meets the core of A_2 in one point, f must be a meridian of U_2 . But this implies that K is trivial, a contradiction. $\square$

Thus $\chi(F_2) \leq 0$. If $\chi(F_2) = 0$, then F_2 consists of annuli, since F_2 cannot contain a Möbius band component. Then $\ell(\sigma) = 2$.

We regard F_2 as the union of $F \cap U_2$ and the rectangles on A again. Let E be an annulus component of F_2 , and let $\partial E = e_1 \cup e_2$. Let g be a loop on ∂U_2 , which is the union of two arcs of $F \cap A$ and two arcs of $\partial F \cap A_2$ and which meets both components of ∂E . Then either g bounds a disk in E , or g is essential on E . In the latter case, we replace $g \cap e_1$ with $e_1 - \text{Int}(g \cap e_1)$. Then the resulting g bounds a disk in E . Since g meets the core of A_2 in two point (with the same sign), g must be a meridian of U_2 . But this implies $p = 2$, a contradiction. Thus we have shown that $\chi(F_2) < 0$. $\square$

Proof of Theorem 1.1(1). By Proposition 3.1, $r = pq$. Consider r -Dehn surgery $K(r)$ on K . Then $K(r) = L(p, q) \sharp L(q, p)$ by [10]. We remark that $L(q, p)$ cannot contain a closed non-orientable surface since $H_1(L(q, p))$ is odd torsion [2]. Let $\widehat{F}$ be the closed non-orientable surface obtained by capping ∂F off by a meridian disk of the attached solid torus of $K(r)$. Then $L(p, q)$ contains a closed non-orientable surface whose genus is $\widehat{F}$ by [2, Corollary 5.2]. Hence $N(p, q) \leq c(K)$.

On the other hand, $N(p, q)$ is also the minimal genus of non-orientable surfaces in a solid torus having a single (p, q) loop as boundary [2]. This means that K bounds a non-orientable surface with genus $N(p, q)$, and thus $c(K) \leq N(p, q)$. Therefore we have $c(K) = N(p, q)$. This completes the proof of Theorem 1.1(1). $\square$

4. THE CASE WHERE K IS ODD

In this section, we prove Theorem 1.1(2).

Suppose that K is odd. Recall that $p > q$. We first determine which of $N(pq - 1, p^2)$ and $N(pq + 1, p^2)$ is smaller. For it, recall the definition of $N(x, y)$ [2]. (Although only the case where x is even is considered in [2], $N(x, y)$ can be defined in general.)

Let x and y be coprime positive integers. We write x/y as a continued fraction:

$$\frac{x}{y} = [a_0, a_1, a_2, \dots, a_n] = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\ddots + \frac{1}{a_n}}}}$$

where the a_i are integers, $a_0 \geq 0$, $a_i > 0$ for $1 \leq i \leq n$, and $a_n > 1$. Note that such an expression is unique (cf. [7]). Then we define b_i inductively as follows:

$$b_0 = a_0,$$

$$b_i = \begin{cases} a_i & \text{if } b_{i-1} \neq a_{i-1} \text{ or if } \sum_{j=0}^{i-1} b_j \text{ is odd,} \\ 0 & \text{if } b_{i-1} = a_{i-1} \text{ and } \sum_{j=0}^{i-1} b_j \text{ is even.} \end{cases}$$

Let $x_i/y_i = [a_0, a_1, a_2, \dots, a_i]$ be the i -th convergent of x/y for $0 \leq i \leq n$. Thus $x/y = x_n/y_n$. Let

$$\sum \left(\frac{x_i}{y_i} \right) = \sum_{j=0}^i b_j,$$

and let

$$N(x, y) = \frac{1}{2} \sum \left(\frac{x_n}{y_n} \right) = \frac{1}{2} \sum \left(\frac{x}{y} \right).$$

That is, $\sum(x/y)$ is obtained by adding the a_i successively except when a partial sum is even we skip the next a_i . [2, Theorem 6.1] showed that $N(x, y)$ gives the minimal genus of closed non-orientable surfaces contained in the lens space $L(x, y)$. (In this case, x must be even. Otherwise, the lens space cannot contain a closed non-orientable surface. Thus $\sum(x/y)$ must be even when x is even.)

Lemma 4.1. *Let $q/p = [a_0, a_1, a_2, \dots, a_n]$, and let b_i ($0 \leq i \leq n$) be defined as above. Then*

$$(pq - 1)/p^2 = \begin{cases} [a_0, a_1, a_2, \dots, a_{n-1}, a_n + 1, a_n - 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1] & \text{if } n \text{ is odd,} \\ [a_0, a_1, a_2, \dots, a_{n-1}, a_n - 1, a_n + 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1] & \text{if } n \text{ is even,} \end{cases}$$

and

$$(pq + 1)/p^2 = \begin{cases} [a_0, a_1, a_2, \dots, a_{n-1}, a_n - 1, a_n + 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1] & \text{if } n \text{ is odd,} \\ [a_0, a_1, a_2, \dots, a_{n-1}, a_n + 1, a_n - 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1] & \text{if } n \text{ is even.} \end{cases}$$

Remark 4.2. In Lemma 4.1, $a_0 = 0$ since $p > q$. Also, if $a_1 = 1$, then $[a_0, \dots, a_3, a_2, a_1]$ should be considered to be $[a_0, \dots, a_3, a_2 + 1]$.

Proof. Since $a_0 = 0$, $p/q = [a_1, a_2, \dots, a_n]$. We show that

$$[a_1, a_2, \dots, a_{n-1}, a_n + 1, a_n - 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1] = \frac{p^2}{pq + (-1)^n}.$$

Let $p_i/q_i = [a_1, \dots, a_i]$ be the i -th convergent of $[a_1, a_2, \dots, a_n]$ ($1 \leq i \leq n$). Then $p_n = p, q_n = q$, and $p_i = a_i p_{i-1} + p_{i-2}$ ($i \geq 3$). By induction, $p_{n-1}/p_{n-2} =$

$[a_{n-1}, a_{n-2}, \dots, a_2, a_1]$. Then

$$\begin{aligned}
[a_n + 1, a_n - 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1] &= [a_n + 1, a_n - 1, [a_{n-1}, a_{n-2}, \dots, a_2, a_1]] \\
&= [a_n + 1, a_n - 1, p_{n-1}/p_{n-2}] \\
&= a_n + 1 + \frac{p_{n-1}}{a_n p_{n-1} - p_{n-1} + p_{n-2}} \\
&= a_n + 1 + \frac{p_{n-1}}{p_n - p_{n-1}} \\
&= \frac{a_n p_n - a_n p_{n-1} + p_n}{p_n - p_{n-1}} \\
&= \frac{a_n p_n + p_{n-2}}{p_n - p_{n-1}}.
\end{aligned}$$

Let us denote this by c_1 . Next, let $c_2 = [a_{n-1}, c_1]$. Then

$$c_2 = \frac{(a_{n-1}a_n + 1)p_n - p_{n-3}}{a_n p_n + p_{n-2}}.$$

Inductively, we define $c_i = [a_{n-i+1}, c_{i-1}]$. For example,

$$\begin{aligned}
c_3 &= [a_{n-2}, c_2] = \frac{(a_{n-2}(a_{n-1}a_n + 1) + a_n)p_n + p_{n-4}}{(a_{n-1}a_n + 1)p_n - p_{n-3}}, \\
c_4 &= [a_{n-3}, c_3] = \frac{(a_{n-3}(a_{n-2}(a_{n-1}a_n + 1) + a_n) + (a_{n-1}a_n + 1))p_n - p_{n-5}}{(a_{n-2}(a_{n-1}a_n + 1) + a_n)p_n + p_{n-4}}.
\end{aligned}$$

We will show $c_n = p^2/(pq + (-1)^n)$.

Let $p'_i/q'_i = [a_n, a_{n-1}, \dots, a_{n-i+1}]$ be the i -th convergent of $[a_n, a_{n-1}, \dots, a_2, a_1]$, $1 \leq i \leq n$. Then $p'_n = p_n = p$, $q'_n = p_{n-1}$, since $p_n/p_{n-1} = [a_n, a_{n-1}, \dots, a_2, a_1]$. Furthermore,

$$\begin{aligned}
p'_1 &= a_n, & p'_2 &= a_{n-1}a_n + 1, & p'_i &= a_{n-i+1}p'_{i-1} + p'_{i-2}, \\
q'_1 &= 1, & q'_2 &= a_{n-1}, & q'_i &= a_{n-i+1}q'_{i-1} + q'_{i-2}.
\end{aligned}$$

Then

$$c_1 = \frac{p'_1 p_n + p_{n-2}}{p_n - p_{n-1}}, \quad c_2 = \frac{p'_2 p_n - p_{n-3}}{p'_1 p_n + p_{n-2}}, \dots, c_{n-2} = \frac{p'_{n-2} p_n + (-1)^{n-1} p_1}{p'_{n-3} p_n + (-1)^n p_2}.$$

Thus

$$c_{n-1} = [a_2, c_{n-2}] = \frac{p'_{n-1} p_n + (-1)^n}{p'_{n-2} p_n + (-1)^{n-1} p_1},$$

since $p_2 = a_1 a_2 + 1$. Finally,

$$c_n = [a_1, c_{n-1}] = \frac{p'_n p_n}{p'_{n-1} p_n + (-1)^n}.$$

Recall that $p'_n = p_n = p$. Also we can show that $p'_n/p'_{n-1} = [a_1, a_2, \dots, a_n] = p/q$ inductively, and so $p'_{n-1} = q$. Therefore $c_n = p^2/(pq + (-1)^n)$ as desired.

Similarly, we can show that

$$[a_1, a_2, \dots, a_{n-1}, a_n - 1, a_n + 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1] = \frac{p^2}{pq + (-1)^{n-1}}.$$

This completes the proof of Lemma 4.1. $\square$

Consider the equation $xq \equiv -1 \pmod{p}$. Since p and q are coprime, this has a solution. In general, if x is a solution, then so is $x + p$. Hence the parity of the solution x is not well-defined, because p is odd. But the equation has the unique solution x satisfying $0 < x < p$. (For, if x and y are such solutions, then $(x-y)q \equiv 0 \pmod{p}$. Then $x \equiv y \pmod{p}$, and so $x = y$.) If such x is even (resp. odd), then the ordered pair (p, q) is said to be of *type A* (resp. *B*).

Let $q/p = [a_0, a_1, a_2, \dots, a_n]$, and let $q_i/p_i = [a_0, a_1, a_2, \dots, a_i]$ be the i -th convergent for $0 \leq i \leq n$. We will use b_i , $0 \leq i \leq n$, defined for $q/p = [a_0, a_1, a_2, \dots, a_n]$ as before.

Lemma 4.3. *If n is even (resp. odd), then the pair (p, q) is of type A if and only if p_{n-1} is even (resp. odd).*

Proof. It is well known that $q_n p_{n-1} - q_{n-1} p_n = (-1)^{n-1}$. Recall $p_n = p, q_n = q$. If n is even, then $p_{n-1} q \equiv -1 \pmod{p}$. Since $0 < p_{n-1} < p_n = p$ ([7]), p_{n-1} gives the unique solution of $xq \equiv -1 \pmod{p}$ such that $0 < x < p$.

If n is odd, then $-p_{n-1} q \equiv -1 \pmod{p}$. Hence $p - p_{n-1}$ gives the unique solution of $xq \equiv -1 \pmod{p}$ such that $0 < x < p$. Since p is odd, p_{n-1} and $p - p_{n-1}$ have distinct parities. $\square$

Remark 4.4. In Lemma 4.3, p_{n-1} and q_{n-1} have distinct parities, and hence when n is odd, (p, q) is of type A if and only if q_{n-1} is even.

Lemma 4.5. *If p_{n-1} is odd, then $b_{n-1} = a_{n-1}$ and $\sum(q_{n-1}/p_{n-1})$ is even.*

Proof. Since p_{n-1} and q_{n-1} have distinct parities, q_{n-1} is even. Let us consider $q_{n-1}/p_{n-1} = [a_0, a_1, \dots, a_{n-1}]$. (If $a_{n-1} = 1$, this is $[a_0, a_1, \dots, a_{n-2} + 1]$.) By [2], $\sum(q_{n-1}/p_{n-1})$ is even, because q_{n-1} is even. (In fact, $\sum(q_{n-1}/p_{n-1})$ is twice the minimal genus of closed non-orientable surfaces contained in the lens space $L(q_{n-1}, p_{n-1})$.)

Now, recall another interpretation of $\sum(q_{n-1}/p_{n-1})$ [2, p107]. Consider the “step” changing $[a_0, a_1, \dots, a_{n-1}] (= [a_0, a_1, \dots, a_{n-2} + a_{n-1}] = [a_0, a_1, \dots, a_{n-2} + 1]$ if $a_{n-1} = 1$) as follows:

$$[a_0, a_1, \dots, a_{n-1}] \rightarrow \begin{cases} [a_0, a_1, \dots, a_{n-1} - 2] & \text{if } a_{n-1} \geq 4, \\ [a_0, a_1, \dots, a_{n-2}, 1] = [a_0, a_1, \dots, a_{n-2} + 1] & \text{if } a_{n-1} = 3, \\ [a_0, a_1, \dots, a_{n-3}] & \text{if } a_{n-1} = 2. \end{cases}$$

Then $\sum(q_{n-1}/p_{n-1})$ is twice the number of steps required to reduce $[a_0, a_1, \dots, a_{n-1}]$ to $[0]$. (Such a reduction works only when q_{n-1} is even.) In other words, the counting process defining $\sum(q_{n-1}/p_{n-1})$ can be done from either end of the sequence

$a_0, a_1, \dots, a_{n-1}$. This implies that the calculation of $\sum(q_{n-1}/p_{n-1})$ involves the last term a_{n-1} . Hence $b_{n-1} = a_{n-1}$. $\square$

Lemma 4.6. *If p_{n-1} is odd, then $N(pq - 1, p^2) = N(pq + 1, p^2) + (-1)^n$.*

Proof. Assume n is odd. Then

$$\begin{aligned} (pq - 1)/p^2 &= [a_0, a_1, a_2, \dots, a_{n-1}, a_n + 1, a_n - 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1], \\ (pq + 1)/p^2 &= [a_0, a_1, a_2, \dots, a_{n-1}, a_n - 1, a_n + 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1] \end{aligned}$$

by Lemma 4.1. In the calculations of $\sum((pq - 1)/p^2)$ and $\sum((pq + 1)/p^2)$, $b_{n-1} = a_{n-1}$ and the partial sum $\sum_{j=0}^{n-1} b_j$ is even by Lemma 4.5. Hence the next term is skipped, that is, $b_n = 0$. Thus we see that $\sum((pq - 1)/p^2) = \sum((pq + 1)/p^2) - 2$. Then $N(pq - 1, p^2) = N(pq + 1, p^2) - 1$.

Assume n is even. Then a similar argument shows that $N(pq - 1, p^2) = N(pq + 1, p^2) + 1$. $\square$

Lemma 4.7. *If p_{n-1} is even, then $N(pq - 1, p^2) + (-1)^n = N(pq + 1, p^2)$.*

Proof. Since $q_{n-1}/p_{n-1} = [a_0, a_1, \dots, a_{n-1}] = [0, a_1, \dots, a_{n-1}] = 1/[a_1, a_2, \dots, a_{n-1}]$, we see $p_{n-1}/q_{n-1} = [a_1, a_2, \dots, a_{n-1}]$. Thus $\sum(p_{n-1}/q_{n-1})$ is even and its calculation involves the last term a_{n-1} , since p_{n-1} is even, as in the proof of Lemma 4.5.

Assume n is odd. Recall

$$\begin{aligned} (pq - 1)/p^2 &= [a_0, a_1, a_2, \dots, a_{n-1}, a_n + 1, a_n - 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1], \\ (pq + 1)/p^2 &= [a_0, a_1, a_2, \dots, a_{n-1}, a_n - 1, a_n + 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1]. \end{aligned}$$

For the calculations of $\sum((pq - 1)/p^2)$ and $\sum((pq + 1)/p^2)$, we use the counting procedure backward as stated in the proof of Lemma 4.5. Starting from a_1 , it reaches $[a_0, a_1, \dots, a_{n-1}, a_n \pm 1, a_n \mp 1, 0] = [a_0, a_1, \dots, a_{n-1}, a_n \pm 1]$. Thus we see that $\sum((pq - 1)/p^2) - 2 = \sum((pq + 1)/p^2)$. Then $N(pq - 1, p^2) - 1 = N(pq + 1, p^2)$.

When n is even, a similar argument shows $N(pq - 1, p^2) + 1 = N(pq + 1, p^2)$. $\square$

We are ready to determine which of $N(pq - 1, p^2)$ and $N(pq + 1, p^2)$ is smaller.

Proposition 4.8. (1) *If the pair (p, q) is of type A, then $N(pq - 1, p^2) + 1 = N(pq + 1, p^2)$.*

(2) *If the pair (p, q) is of type B, then $N(pq + 1, p^2) + 1 = N(pq - 1, p^2)$.*

Proof. Suppose (p, q) is of type A. If n is even, then so is p_{n-1} by Lemma 4.3. Then $N(pq - 1, p^2) + 1 = N(pq + 1, p^2)$ by Lemma 4.7. If n is odd, then p_{n-1} is odd. Then $N(pq - 1, p^2) = N(pq + 1, p^2) - 1$ by Lemma 4.6. This proves (1). (2) can be proved similarly. $\square$




FIGURE 4.

Remark 4.9. In general, the difference between $N(x, y)$ and $N(x+2, y)$ can be big. For example, $N(26, 25) = 13$, $N(28, 25) = 6$.

Now, we will explain the geometric meaning of types A and B. Recall that $K = T(p, q)$ lies on the standard torus T , which decomposes S^3 into two solid tori V and W , and that K runs p times longitudinally and q times meridionally with respect to V . Choose an arc γ on T as shown in Figure 4. (Here, the end circles of the cylinder are identified to form T .) Then $\partial\gamma$ splits K into two arcs K_a and K_b . Let $K_A = \gamma \cup K_a$ and $K_B = \gamma \cup K_b$ as shown in Figure 4. Clearly, both K_A and K_B are torus knots, which are uniquely determined by K . In Figure 4, $K = T(5, 3)$, $K_A = T(3, 2)$ and $K_B = T(2, 1)$ (with respect to V).

Let $K_A = T(r_1, s_1)$ and $K_B = T(r_2, s_2)$. Then $p = r_1 + r_2$ and $q = s_1 + s_2$.

Lemma 4.10. (1) *If (p, q) is of type A, then r_1 and s_2 are even, and hence s_1 and r_2 are odd.*

(2) *If (p, q) is of type B, then s_1 and r_2 are even, and hence r_1 and s_2 are odd.*

Proof. Let m be a meridian of V near γ , and label the points of $K \cap m$, $0, 1, 2, \dots, p-1$ along m as shown in Figure 4. Start the point with label 0 and follow K_A in the direction of γ . Then we will come back to the point with label $1 + q$ after running once longitudinally. Hence K_A gives the equation $1 + r_1 q \equiv 0 \pmod{p}$. Thus if (p, q) is of type A, then r_1 is even, and s_1 is odd. Since $p = r_1 + r_2$ and $q = s_1 + s_2$ are odd, r_2 is odd and s_2 is even. This proves (1). (2) follows similarly. $\square$

Lemma 4.11. (1) *If (p, q) is of type A, then K bounds a non-orientable surface with genus $N(r_1, s_1) + N(s_2, r_2)$ and boundary slope $pq - 1$.*




FIGURE 5.

- (2) If (p, q) is of type B, then K bounds a non-orientable surface with genus $N(s_1, r_1) + N(r_2, s_2)$ and boundary slope $pq + 1$.

Proof. Assume that (p, q) is of type A. By Lemma 4.10, K_A bounds a non-orientable surface F_A contained in V with genus $N(r_1, s_1)$, and K_B bounds such F_B in W with genus $N(s_2, r_2)$. Let $G = F_A \cup F_B$. Since $F_A \cap F_B = \gamma$, $\partial G = K$ and G has genus $N(r_1, s_1) + N(s_2, r_2)$. It is easy to see that G has the desired boundary slope from Figure 5. This proves (1).

A similar argument shows (2). $\square$

Recall that $q_{n-1}/p_{n-1} = [a_0, a_1, \dots, a_{n-1}]$ is the $(n-1)$ th convergent of $q_n/p_n = q/p = [a_0, a_1, \dots, a_n]$.

Lemma 4.12. *If n is even, then $r_1 = p_{n-1}$ and $s_1 = q_{n-1}$. If n is odd, $r_2 = p_{n-1}$ and $s_2 = q_{n-1}$.*

Proof. Suppose n is even. As shown in the proof of Lemma 4.3, we see $r_1 = p_{n-1}$. Recall that $q_{n-1}p - p_{n-1}q = 1$ and $0 < q_{n-1} < q_n = q$. By considering the intersection number between K and K_A on T (after giving K an orientation, which induces that of K_A), we have $|s_1p - p_{n-1}q| = 1$. Since $p \geq 5$, we have $(q_{n-1} - s_1)p = 0$, and then $s_1 = q_{n-1}$.

When n is odd, we see $r_1 = p - p_{n-1}$. Then a similar argument shows that $s_1 = q - q_{n-1}$. $\square$

Lemma 4.13. $[a_1, a_2, \dots, a_{n-1}, a_n - 1] = (p - p_{n-1})/(q - q_{n-1})$.

Proof. Recall that $p_{n-1}/q_{n-1} = [a_1, a_2, \dots, a_{n-1}]$. Put $P/Q = [a_1, a_2, \dots, a_{n-1}, a_n - 1]$. If $a_n > 2$, then $P = (a_n - 1)p_{n-1} + p_{n-2} = p - p_{n-1}$ as desired. If $a_n = 2$, then $P/Q = [a_1, a_2, \dots, a_{n-1} + 1]$. Thus $P = (a_{n-1} + 1)p_{n-2} + p_{n-3} = p_{n-1} + p_{n-2}$. Since $p = p_n = a_n p_{n-1} + p_{n-2} = 2p_{n-1} + p_{n-2}$, $p_{n-1} + p_{n-2} = p - p_{n-1}$ as desired again. Similarly for Q . $\square$

Lemma 4.14. (1) If (p, q) is of type A, then $N(pq - 1, p^2) = N(r_1, s_1) + N(s_2, r_2)$.

(2) If (p, q) is of type B, then $N(pq + 1, p^2) = N(s_1, r_1) + N(r_2, s_2)$.

Proof. (1) Suppose that p_{n-1} is odd. By Lemma 4.3, n is odd. Then $K_B = T(p_{n-1}, q_{n-1})$ by Lemma 4.12, and hence $K_A = T(p - p_{n-1}, q - q_{n-1})$. Recall that $(pq - 1)/p^2 = [a_0, a_1, a_2, \dots, a_{n-1}, a_n + 1, a_n - 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1]$ by Lemma 4.1. In the counting procedure of $\sum((pq - 1)/p^2)$, $\sum_{j=0}^{n-1} b_j = \sum(q_{n-1}/p_{n-1})$ by Lemma 4.5.

As stated in the proof of Lemma 4.5, the counting procedure can be done backward. But $[a_1, a_2, \dots, a_{n-1}, a_n - 1] = (p - p_{n-1})/(q - q_{n-1})$ by Lemma 4.13. Thus

$$\sum((pq - 1)/p^2) = \sum(q_{n-1}/p_{n-1}) + \sum((p - p_{n-1})/(q - q_{n-1})).$$

Hence we have $N(pq - 1, p^2) = N(s_2, r_2) + N(r_1, s_1)$.

Suppose that p_{n-1} is even. Then n is even by Lemma 4.3 so that $K_A = T(p_{n-1}, q_{n-1})$. Recall $(pq - 1)/p^2 = [a_0, a_1, a_2, \dots, a_{n-1}, a_n - 1, a_n + 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1]$. Perform the counting of $\sum((pq - 1)/p^2)$ backward. Then $\sum[a_1, a_2, \dots, a_{n-1}] = \sum(p_{n-1}/q_{n-1})$, and $\sum[a_0, a_1, \dots, a_{n-1}, a_n - 1] = \sum((q - q_{n-1})/(p - p_{n-1}))$. Hence $N(pq - 1, p^2) = N(s_2, r_2) + N(r_1, s_1)$ again.

A similar argument shows (2). $\square$

Corollary 4.15. $c(K) \leq N(pq - 1, p^2)$ (resp. $N(pq + 1, p^2)$) when K is of type A (resp. B).

Proof. By Lemmas 4.11 and 4.14, if (p, q) is of type A (resp. B), then K can bound a non-orientable surface with genus $N(pq - 1, p^2)$ (resp. $N(pq + 1, p^2)$). Hence $c(K) \leq N(pq - 1, p^2)$ (resp. $N(pq + 1, p^2)$) when K is of type A (resp. B). $\square$

We now determine the boundary slope r of a minimal genus non-orientable spanning surface F of K .

Proposition 4.16. $r = pq \pm 1$.

Proof. Suppose not. Since $\Delta = |r - pq|$ is odd, $\Delta \geq 3$. The proof is divided into two cases.

Case 1. $\Delta \geq 5$.

We perform $(\Delta - 1)/2$ disk splittings for F along mutually disjoint $(\Delta - 1)/2$ rectangles on A as in the proof of Proposition 3.1. By this operation, ∂F breaks up into a loop with slope $pq \pm 1$ and the others inessential on $\partial E(K)$. It can be observed that the latter is not empty under the condition $\Delta \geq 5$. Let F_1 be the resulting connected surface whose boundary contains a component of slope $pq \pm 1$, and let F_2 be the others. Note that $\chi(F) = \chi(F_1) + \chi(F_2)$. Thus F_1 is non-orientable, and F_2 may be empty or disconnected. Also $F_1 \cap A \neq \emptyset$, but $F_2 \cap A = \emptyset$.

Now, A splits $E(K)$ into two solid tori U_1, U_2 , where we assume $F_2 \subset U_2$. Let $A_i = \partial E(K) \cap U_i$. Define σ by using the arcs $\partial F \cap A_2$ as in the proof of Proposition 3.1.

Claim 4.17. $F_2 \neq \emptyset$ and F_1 has a single boundary component.




FIGURE 6.




FIGURE 7.

Proof of Claim 4.17. The argument in the proof of Claim 3.3 works, and so F_2 has no disk components. Thus $\chi(F) = \chi(F_1) + \chi(F_2) \leq \chi(F_1)$. If either $F_2 = \emptyset$ or $|\partial F_1| > 1$, then ∂F_1 contains an inessential component on $\partial E(K)$. Then cap all inessential components of ∂F_1 off by disks on $\partial E(K)$. The resulting surface gives a non-orientable spanning surface of K with fewer betti number than F . This contradicts the minimality of F . $\square$

Recall that $\ell(\sigma)$ is the (common) length of the orbits of σ .

Claim 4.18. $\ell(\sigma) = 1$.

Proof of Claim 4.18. Suppose not. Let α be the arc of $F \cap A$ which is disjoint from all rectangles used in the disk splittings. This arc remains in F_1 . We choose the labeling so that the point $\alpha \cap a_1$ has Δ . Since $\ell(\sigma) \neq 1$, the point $\alpha \cap a_1$ is not connected with $\alpha \cap a_2$ (with label Δ) by an arc in A_2 . See Figure 6 where $\Delta = 7$. (In this and successive figures, we assume $r = pq + \Delta$. But the situation when $r = pq - \Delta$ is similar.)

If either situation as shown in Figure 7 happens, then the disk splittings give rise to inessential components on both A_1 and A_2 . Then ∂F_1 would have an inessential component, which is impossible by Claim 4.17. Thus the only possible configurations are as shown in Figure 8. But it is easy to see that ∂F_1 has an inessential component (and $F_2 = \emptyset$) in either case. This contradicts Claim 4.17 again. $\square$

Thus $|\partial F_2| = (\Delta - 1)/2 \geq 2$. If $\chi(F_2) = 0$, then F_2 consists of annuli. Then the same argument as in the proof of Claim 3.3 gives a contradiction. (That is, let f be a loop consisting of an arc of $F \cap A$ and one of $\partial F \cap A_2$, which meets a component E of F_2 . Then either f bounds a disk in E , or f is essential in E . In



FIGURE 8.




FIGURE 9.

the latter, replace $f \cap e$ with $e - \text{Int}(f \cap e)$, where e is the component of ∂E meeting f . Then it is necessarily inessential in E .) Thus we have shown that $\chi(F_2) < 0$, and so $\chi(F) < \chi(F_1)$, a contradiction.

Case 2. $\Delta = 3$.

We perform a disk splitting for F along one rectangle on A as before. Use the same notation as in Case 1. Then $F_2 = \emptyset$ or $|\partial F_2| = 1$. In the latter case, the argument in the proof of Claim 3.3 shows that F_2 is not a disk. Thus $\chi(F_2) < 0$. Then $\chi(F) < \chi(F_1)$, a contradiction. Hence $F_2 = \emptyset$, and the only possible configurations are as shown in Figure 9.

Claim 4.19. *If $r = pq + 3$ (resp. $pq - 3$), then (p, q) is of type A (resp B).*

Proof of Claim 4.19. Suppose that $r = pq + 3$ and that (p, q) is of type B. Let F' be the resulting surface obtained from F by a disk splitting. Then F' has boundary slope $pq - 1$, and $\chi(F') = \chi(F)$. Note that F' is connected and non-orientable. Hence $N(pq - 1, p^2) \leq c(K)$. This contradicts that $c(K) \leq N(pq + 1, p^2) < N(pq - 1, p^2)$ (Proposition 4.8 and Corollary 4.15).

The case where $r = pq - 3$ is similar. □

We will exclude $r = pq + 3$. The argument to the case $r = pq - 3$ is similar.

Recall $E(K) = U_1 \cup_A U_2$ and $A_i = \partial E(K) \cap U_i$. We assume that K has type (p, q) with respect to U_1 . Let $G_i = F \cap U_i$ and $K_i = \partial G_i$ for $i = 1, 2$. Then $\chi(G_1) + \chi(G_2) - 3 = \chi(F)$, and so $\beta_1(G_1) + \beta_1(G_2) + 2 = \beta_1(F)$.

Let us also recall that $q/p = [a_0, a_1, \dots, a_n]$ and $q_{n-1}/p_{n-1} = [a_0, a_1, \dots, a_{n-1}]$.

- Claim 4.20.** (1) If n is odd, K_1 has type $(3p_{n-1} + p(3k - 1), 3q_{n-1} + q(3k - 1))$ with respect to U_1 , and K_2 has type $(-3q_{n-1} + q(2 - 3k), -3p_{n-1} + p(2 - 3k))$ with respect to U_2 , for some even k .
- (2) If n is even, K_1 has type $(-3p_{n-1} + p(3k - 1), -3q_{n-1} + q(3k - 1))$ with respect to U_1 , and K_2 has type $(3q_{n-1} + q(2 - 3k), 3p_{n-1} + p(2 - 3k))$ with respect to U_2 , for some odd k .

Proof of Claim 4.20. (1) Let M denote the matrix

$$\begin{pmatrix} p_{n-1} & q_{n-1} \\ p & q \end{pmatrix}.$$

Note $\det M = (-1)^{n-1} = 1$. Both A_i are parallel to the annulus $T - \text{Int}A$ in $N(K)$. Thus K_i is isotopic to K'_i , say, on T rel A . Let f be an orientation-preserving self-homeomorphism on T corresponding to M . Then it is easy to see from Figure 9 that $f^{-1}(K'_1)$ has type $(3, 3k - 1)$ and $f^{-1}(K'_2)$ has type $(-3, 2 - 3k)$ for some integer k (with suitable orientations). The conclusion follows easily from this.

Remark that p_{n-1} is odd and q_{n-1} is even by Claim 4.19, Lemma 4.3 and the remark following it. Since K_i bounds G_i in U_i , we see that k is even.

(2) Use the matrix

$$\begin{pmatrix} -p_{n-1} & -q_{n-1} \\ p & q \end{pmatrix}$$

instead of M . □

Claim 4.21. (1) Let n be odd. Then

$$\frac{3p_{n-1} + p(3k - 1)}{3q_{n-1} + q(3k - 1)} = \begin{cases} [a_1, a_2, \dots, a_{n-1}, a_n - 1, 1, |k| - 1, 3] & \text{if } k \leq -2, \\ [a_1, a_2, \dots, a_{n-1}, a_n - 3] & \text{if } k = 0, \\ [a_1, a_2, \dots, a_{n-1}, a_n, k - 1, 1, 2] & \text{if } k \geq 2 \end{cases}$$

and

$$\frac{-3q_{n-1} + q(2 - 3k)}{-3p_{n-1} + p(2 - 3k)} = \begin{cases} [0, a_1, a_2, \dots, a_{n-1}, a_n - 1, 1, |k| - 1, 1, 2] & \text{if } k \leq -2, \\ [0, a_1, a_2, \dots, a_{n-1}, a_n - 2, 2] & \text{if } k = 0, \\ [0, a_1, a_2, \dots, a_{n-1}, a_n, k - 1, 3] & \text{if } k \geq 2. \end{cases}$$

(2) Let n be even. Then

$$\frac{-3p_{n-1} + p(3k - 1)}{-3q_{n-1} + q(3k - 1)} = \begin{cases} [a_1, a_2, \dots, a_{n-1}, a_n, |k|, 3] & \text{if } k \leq -1, \\ [a_1, a_2, \dots, a_{n-1}, a_n - 2, 2] & \text{if } k = 1, \\ [a_1, a_2, \dots, a_{n-1}, a_n - 1, 1, k - 2, 1, 2] & \text{if } k \geq 3 \end{cases}$$

and

$$\frac{3q_{n-1} + q(2-3k)}{3p_{n-1} + p(2-3k)} = \begin{cases} [0, a_1, a_2, \dots, a_{n-1}, a_n, |k|, 1, 2] & \text{if } k \leq -1, \\ [0, a_1, a_2, \dots, a_{n-1}, a_n - 3] & \text{if } k = 1, \\ [0, a_1, a_2, \dots, a_{n-1}, a_n - 1, 1, k - 2, 3] & \text{if } k \geq 3. \end{cases}$$

(We consider that $[\dots, a_{n-1}, -1] = [\dots, a_{n-1} - 1]$, $[\dots, a_{n-2}, a_{n-1}, 0] = [\dots, a_{n-2}]$, and $[\dots, a_{n-1}, 0, 2] = [\dots, a_{n-1} + 2]$.)

Proof of Claim 4.21. This is straightforward. Suppose that n is odd and $k \leq -2$. Then

$$\begin{aligned} [a_1, a_2, \dots, a_{n-1}, a_n - 1, 1, |k| - 1, 3] &= [a_1, a_2, \dots, a_{n-1}, [a_n - 1, 1, |k| - 1, 3]] \\ &= \frac{[a_n - 1, 1, |k| - 1, 3]p_{n-1} + p_{n-2}}{[a_n - 1, 1, |k| - 1, 3]q_{n-1} + q_{n-2}} \\ &= \frac{(a_n - 1)p_{n-1} + p_{n-2} + \frac{3k+2}{3k-1}p_{n-1}}{(a_n - 1)q_{n-1} + q_{n-2} + \frac{3k+2}{3k-1}q_{n-1}} \\ &= \frac{p - p_{n-1} + \frac{3k+2}{3k-1}p_{n-1}}{q - q_{n-1} + \frac{3k+2}{3k-1}q_{n-1}} \\ &= \frac{(3k-1)p + 3p_{n-1}}{(3k-1)q + 3q_{n-1}} \end{aligned}$$

as desired. The other cases can be checked similarly. $\square$

By using these continued fractions, we can evaluate $\beta_1(G_i)$. Assume n is odd. As in the proof of Lemma 4.14,

$$\beta_1(G_1) \geq N(3p_{n-1} + p(3k-1), 3q_{n-1} + q(3k-1)) = \begin{cases} N(p - p_{n-1}, q - q_{n-1}) - 1 & \text{if } k = 0, \\ N(p - p_{n-1}, q - q_{n-1}) + \frac{|k|}{2} + 1 & \text{otherwise,} \end{cases}$$

and

$$\beta_1(G_2) \geq N(-3q_{n-1} + q(2-3k), -3p_{n-1} + p(2-3k)) = \begin{cases} N(q_{n-1}, p_{n-1}) + 1 & \text{if } k = 0, \\ N(q_{n-1}, p_{n-1}) + \frac{|k|}{2} + 1 & \text{otherwise.} \end{cases}$$

Recall that $\beta_1(F) = \beta_1(G_1) + \beta_1(G_2) + 2$. Thus $c(K) = \beta_1(F) > N(pq - 1, p^2)$, which contradicts Corollary 4.15.

The argument to the case n even is similar. $\square$

Proposition 4.22. *If (p, q) is of type A (resp. B), then $r = pq - 1$ (resp. $pq + 1$).*

Proof. By Proposition 4.16, $r = pq - 1$ or $pq + 1$. Suppose that (p, q) is of type A. If $r = pq + 1$, then $pq + 1$ -surgery $K(pq + 1) = L(pq + 1, p^2)$ contains the closed non-orientable surface $\widehat{F}$, obtained by capping ∂F off by a meridian disk of the

attached solid torus. Hence $N(pq+1, p^2) \leq c(K)$. This contradicts Lemma 4.8 and Corollary 4.15. Thus $r = pq - 1$.

A similar argument shows that $r = pq + 1$ when (p, q) is of type B. $\square$

Proof of Theorem 1.1(2). Assume (p, q) is of type A. If F is a non-orientable spanning surface of K realizing $c(K)$, then its boundary slope r is $pq - 1$ by Proposition 4.22. We can cap ∂F off by a meridian disk of the attached solid torus of $K(r)$. Thus $N(pq-1, p^2) \leq c(K)$. Combined with Corollary 4.15, we have $c(K) = N(pq-1, p^2)$. Similarly, if (p, q) is of type B, then $c(K) = N(pq+1, p^2)$. This completes the proof of Theorem 1.1(2). $\square$

Proof of Theorem 1.2. By [11, Proposition 4.3], $c(K_1 \# K_2) = c(K_1) + c(K_2)$ if and only if $c(K_1) = \Gamma(K_1)$ and $c(K_2) = \Gamma(K_2)$, where $\Gamma(K_i) = \min\{2g(K_i), c(K_i)\}$. Let $K = T(p, q)$ be a non-trivial torus knot. As known well, $g(K) = (p-1)(q-1)/2$. As in the proof of Lemma 2.2, $c(K) \leq \min\{(p-1)q/2, (q-1)p/2\}$. Since it is easy to check that $\min\{(p-1)q/2, (q-1)p/2\} < 2g(K)$, we have $\Gamma(K) = c(K)$. Thus, if K_1 and K_2 are torus knots, then $c(K_1 \# K_2) = c(K_1) + c(K_2)$.

By a standard cut-and-paste argument, $\Gamma(K_1 \# K_2) = \Gamma(K_1) + \Gamma(K_2)$. (See also [11, Section 0].) Thus we have the result inductively. $\square$

5. EXAMPLES AND COMMENTS

Let K be a torus knot, and let F be a minimal genus non-orientable surface bounded by K .

Example 5.1. Let $K = T(8, 3)$. Since K is even, the boundary slope of F is 24. We have $8/3 = [2, 1, 2]$, so that $b_0 = 2, b_1 = 0, b_2 = 2$. This gives $\sum(8/3) = 4$, and hence $c(K) = N(8, 3) = 2$.

Example 5.2. Let $K = T(7, 5)$. Since $4 \cdot 5 \equiv -1 \pmod{7}$, the pair $(7, 5)$ is of type A. Thus the boundary slope of F is 34. Since $34/49 = [0, 1, 2, 3, 1, 3]$, $c(K) = N(34/49) = (0 + 2 + 1 + 3)/2 = 3$.

Next, let $K = T(25, 9)$. Then $(25, 9)$ is of type B. Then the boundary slope of F is 226. Since $226/625 = [0, 2, 1, 3, 3, 1, 3, 1, 2]$, $c(K) = N(226/625) = (0 + 1 + 3 + 1 + 3 + 2)/2 = 5$.

In this paper, we proved that the boundary slope of a minimal genus non-orientable spanning surface for a non-trivial torus knot is unique. It seems that such a surface itself is unique up to isotopy in the knot exterior.

REFERENCES

1. K. Bessho, *Incompressible surfaces bounded by links*, Master thesis (in Japanese), Osaka University, 1994.
2. G. Bredon and J. Wood, *Non-orientable surfaces in orientable 3-manifolds*, Invent. Math. **7** (1969), 83–110.

3. B. E. Clark, *Crosscaps and knots*, Internat. J. Math. and Math. Sci. **1** (1978), 113–123.
4. W. End, *Non-orientable surfaces in 3-manifolds*, Arch. Math. **59** (1992), 173–185.
5. C. Frohman, *One-sided incompressible surfaces in Seifert fibered spaces*, Topology Appl. **23** (1986), 103–116.
6. C. McA. Gordon, *Dehn surgery and satellite knots*, Trans. Amer. Math. Soc. **275** (1983), 687–708.
7. G. Hardy and E. Wright, *An introduction to the theory of numbers, Fifth edition*, The Clarendon Press, Oxford University Press, New York, 1979.
8. K. Ichihara, M. Ohtouge and M. Teragaito, *Boundary slopes of non-orientable Seifert surfaces for knots*, Topology Appl. **122** (2002), 467–478.
9. W. Jaco and P. Shalen, *Seifert fibered spaces in 3-manifolds*, Mem. Amer. Math. Soc. **21** (1979), no.220.
10. L. Moser, *Elementary surgery along a torus knot*, Pacific J. Math. **38** (1971), 737–745.
11. H. Murakami and A. Yasuhara, *Crosscap number of a knot*, Pacific J. Math. **171** (1995), 261–273.
12. ———, *Four-genus and four-dimensional clasp number of a knot*, Proc. Amer. Math. Soc. **128** (2000), 3693–3699.
13. D. Rolfsen, *Knots and Links*, Math. Lect. Ser. **7**, Publish or Perish, Berkeley, California, 1976.
14. J. H. Rubinstein, *One-sided Heegaard splittings of 3-manifolds*, Pacific J. Math. **76** (1978), 185–200.
15. M. Teragaito, *Creating Klein bottles by surgery on knots*, J. Knot Theory Ramifications **10** (2001), 781–794.
16. ———, *Dehn surgeries on composite knots creating Klein bottles*, J. Knot Theory Ramifications **8** (1999), 391–395.
17. C. M. Tsau, *A note on incompressible surfaces in solid tori and in lens spaces*, in Knots 90 (Osaka, 1990), 213–229, de Gruyter, Berlin, 1992.
18. J. Uhing, *Zur Kreuzhaubenzahl von Knoten*, diploma thesis, Universitat Dortmund.

DEPARTMENT OF MATHEMATICS AND MATHEMATICS EDUCATION, FACULTY OF EDUCATION, HIROSHIMA UNIVERSITY, 1-1-1 KAGAMIYAMA, HIGASHI-HIROSHIMA 739-8524, JAPAN
E-mail address: `teragai@hiroshima-u.ac.jp`